

ACKNOWLEDGEMENT

This work was supported by the by the Italian Ministry of Education, University and Scientific Research (MIUR) under the project SIR “SNOWAVE” RBSI148WE5.

The authors would like to thank Regione Valle d’Aosta and Fondazione Montagna Sicura for the support during the experimental campaigns, and the Finnish Meteorological Institute for providing the snow fork.

REFERENCES

- [1] Vaughan, D. G., et al: Observations: Cryosphere, Climate change 2013: The physical science basis, Cambridge University Press, 2013.
- [2] Baggi, S. and Schweizer, J.: Characteristics of wet-snow avalanche activity: 20 years of observations from a high alpine valley (Dischma, Switzerland), *Natur. Hazards*, 50(1), 97–108, 2009.
- [3] Bacchi, B. and Ranzi, R.: Hydrological and meteorological aspects of floods in the Alps: an overview, *Hydrology and Earth System Sciences*, 7(6), 784–798, 2003.
- [4] Bartelt, P. and Lehning, M.: A physical SNOWPACK model for the Swiss avalanche warning – Part I: numerical model. *Cold Regions Science and Technology*, 35(23), 2002.
- [5] Heilig, A., Eisen, O., and Schneebeli, M.: Temporal observations of a seasonal snowpack using upward-looking GPR, *Hydrol. Processes*, 24(22), 3133–3145, 2010.
- [6] Okorn, R., et al.: Upward-looking L-band FMCW radar for snow cover monitoring, *Cold Regions Sci. Technol.*, 103, 31–40, 2014.
- [7] Godio, A., et al.: Seasonal monitoring of snow properties by WCR and up-GPR, 21st Eur. Meeting Environ. Eng. Geophys. Near Surf. Geosci., Turin, Italy, September 6–10, 2015.
- [8] Schmid, L., et al.: A novel sensor combination (upGPR-GPS) to continuously and nondestructively derive snow cover properties, *Geophys. Res. Lett.*, 42(9), 3397–3405, 2015.
- [9] Bradford, J. H., Harper, J. T., and Brown, J.: Complex dielectric permittivity measurements from ground-penetrating radar data to estimate snow liquid water content in the pendular regime, *Water Resource Research*, 45, 2009.
- [10] Pasion, M., Barbolini, M., Dell’Acqua, F., Espín-López, P. F., and Silvestri, L.: Snowpack Monitoring Using a Dual-Receiver Radar Architecture, *IEEE Transactions on Geoscience and Remote Sensing*, 57(2), 1195–1204, 2019.
- [11] Barbolini, M., Dell’Acqua, F., Espín-López, P. F., Silvestri, L., and Pasion, M.: Estimate of snowpack parameters for dry and wet snow using a dual-receiver, stand-alone FMCW radar architecture, *Cold Regions Science and Technology*, submitted.
- [12] Espín-López, P. F., Pasion, M., Alaimo, G., Marconi, S., Auricchio, F., Heinänen, V., and Järveläinen, J.: 3-D Printed Antenna for Snowpack Monitoring, *IEEE Antennas and Wireless Propagation Letters*, 17(11), 2109–2113, 2018.
- [13] Hallikainen, M. T., Ulaby, F. T., and Abdelrazik, M.: Dielectric properties of snow in the 3 to 37 GHz range, *IEEE Transactions on Antennas and Propagation*, 34(11), 1329–1340, 1986.
- [14] Sihvola, A. H. and Tiuri, M. E.: Snow fork for field determination of the density and wetness profiles of a snow pack, *IEEE Transactions on Geoscience and Remote Sensing*, 24(5), 717–721, 1986.